

# Insecticidal efficacy of natural diatomaceous earth deposits from Greece and Romania against four stored grain beetles: the effect of temperature and relative humidity

Christos G. ATHANASSIOU<sup>1</sup>, Nickolas G. KAVALLIERATOS<sup>2,3</sup>, Andrei CHIRILOAIE<sup>4</sup>, Thomas N. VASSILAKOS<sup>1</sup>, Viorel FĂTU<sup>4</sup>, Sonica DROSU<sup>4</sup>, Maria CIOBANU<sup>4</sup>, Roxana DUDOIU<sup>4</sup>

<sup>1</sup>Department of Agriculture Crop Production and Rural Environment, University of Thessaly, Greece

<sup>2</sup>Department of Crop Science, Agricultural University of Athens, Greece

<sup>3</sup>Department of Entomology and Agricultural Zoology, Benaki Phytopathological Institute, Kifissia, Greece

<sup>4</sup>Research Development Institute for Plant Protection Bucharest, Romania

## Abstract

In two separate experimental bioassays, the efficacy of diatomaceous earth (DE) deposits from Greece and Romania, as well as two DE commercial formulations were evaluated against four stored-product beetle species. In the 1<sup>st</sup> experiment, four doses (100, 300, 500 and 900 ppm) of two Greek (Elassona 1 and Elassona 2), two Romanian (PatRom and AdRom) DEs and two commercial DEs (SilicoSec and PyriSec) were tested against adults of *Tribolium castaneum* (Herbst) (Coleoptera Tenebrionidae) and *Oryzaephilus surinamensis* (L.) (Coleoptera Silvanidae) in wheat and barley. In the 2<sup>nd</sup> experiment, the effect of temperature (20, 25 and 30 °C) and relative humidity (RH) (55 and 75%) on the efficacy of DEs from Greece (Elassona 1), Romania (PatRom) and the commercial DE SilicoSec at three doses (300, 600 and 900 ppm) in wheat was investigated against adults of *Sitophilus oryzae* (L.) (Coleoptera Curculionidae) and *Rhyzopertha dominica* (F.) (Coleoptera Bostrychidae). Mortality of the exposed adults in both experiments was assessed after 7, 14 and 21 days of exposure. In the 1<sup>st</sup> experiment, the most effective DE was PyriSec which caused complete mortality of *O. surinamensis* adults at 300 ppm after 7 and 14 days of exposure in wheat and barley respectively. In both commodities, the Greek (Elassona 1) and Romanian (PatRom) DEs had the same performance with SilicoSec at 500 ppm. Complete mortality of *T. castaneum* adults was recorded after 21 days at the highest dose in both commodities treated with PyriSec. Greek and Romanian DEs caused low mortality (< 25%) even after 21 days of exposure at the highest dose. *T. castaneum* was less susceptible than *O. surinamensis*, especially to Greek and Romanian DEs. Results from the 2<sup>nd</sup> experiment indicated that *S. oryzae* adults were more susceptible than *R. dominica* adults to the tested DEs. Mortality levels for both species were high, but only at 900 ppm. The increase of temperature, increased the efficacy of the tested DEs against adults of both species. Higher mortality levels were recorded in the low RH for both species. The Romanian DE PatRom was more effective compared with SilicoSec and the Greek DE Elassona 1 against *S. oryzae* and *R. dominica* adults at 900 ppm and 25 °C.

**Key words:** diatomaceous earths, *Rhyzopertha dominica*, *Sitophilus oryzae*, *Tribolium castaneum*, *Oryzaephilus surinamensis*, temperature, relative humidity.

## Introduction

The rice weevil, *Sitophilus oryzae* (L.) (Coleoptera Curculionidae), the lesser grain borer, *Rhyzopertha dominica* (Coleoptera Bostrychidae), the red flour beetle, *Tribolium castaneum* (Herbst) (Coleoptera Tenebrionidae), and the saw toothed grain beetle, *Oryzaephilus surinamensis* (L.) (Coleoptera Silvanidae) are destructive cosmopolitan pests of a wide variety of stored products (Hill, 1990). Several chemical insecticides are used for their control, but the future of some of them remains uncertain, due to both health and environmental concerns (Arthur, 2012). Diatomaceous earths (DEs) are used as alternatives to chemical insecticides and fumigants due to certain advantages; they exhibit low mammalian toxicity, their origin is natural (fossils of phytoplanktons) and they have non-chemical mode of action by absorbing the waterproof epicuticular lipids from insects' cuticle which die because of desiccation (Ebeling, 1971; Korunic, 1998). Several previous studies have demonstrated that DEs are very effective against stored-product insects (Fields and Korunic, 2000; Arthur, 2000a; 2000b; Subramanyam and Roesli, 2000; Arthur, 2002; Arthur and Throne, 2003; Athanassiou *et al.*, 2003; 2004; 2005; 2011, 2014; Andrić *et al.*, 2012). The efficacy of DEs

against stored grain pests is influenced by various factors, i.e., insect species, strain or stage, geographical origin, commodity, temperature and relative humidity (RH) (Le Patourel, 1986; Desmarchelier and Dines, 1987; Aldryhim, 1993; Golob, 1997; Arthur, 2000b; Stathers *et al.*, 2004; Kavallieratos *et al.*, 2007; Athanassiou *et al.*, 2011; 2014; Baldassari and Martini, 2014). The investigation for the discovery of novel DEs with insecticidal properties enhances the available tools for the Integrated Pest Management of stored-product insects (Athanassiou *et al.*, 2011; Nwaubani *et al.*, 2014). Although previous reports have indicated that DE deposits from southeastern Europe are potentially effective for the control of stored-product insect pests (Vayias *et al.*, 2009; Rojht *et al.*, 2010; Athanassiou *et al.*, 2011; Andrić *et al.*, 2012), very little information is available on the insecticidal efficacy of mined DEs in Romania. The only available studies on this issue by Chiriloaie *et al.* (2013; 2014) have pointed out that DEs originating from Romania can satisfactorily control *S. oryzae* in stored wheat. Therefore, the objective of the present study was to evaluate the insecticidal efficacy of DEs from Romania and compare them with commercially available DE formulations and DEs from Greece against *S. oryzae*, *R. dominica*, *T. castaneum* and *O. surinamensis*. Furthermore, the in-

fluence of temperature, RH and type of commodity in their insecticidal efficacy was also investigated. For this purpose two separate experimental tests were conducted in Romania and Greece.

## Materials and methods

### Insects

Adults of mixed sex, with age < 21 days, reared at 25 °C, 60% RH and continuous darkness were used in the tests. For the 1<sup>st</sup> experiment, adults of *O. surinamensis* and *T. castaneum* were used from laboratory populations that are kept at the Research Development Institute for Plant Protection Bucharest. The *O. surinamensis* individuals were cultured in whole wheat. The *T. castaneum* individuals were cultured in mixture of maize and wheat flour, with 5% brewer yeast. For the 2<sup>nd</sup> experiment, adults of *S. oryzae* and *R. dominica* were used from laboratory cultures that are kept in the Laboratory of Entomology and Agricultural Zoology, University of Thessaly in Greece. Individuals of both species were reared in whole wheat.

### DEs processing and commodity application

The amorphous DE samples from two local DE deposits from Romania, Buzău Valey-Pătărlagele (namely PatRom) and South of Dobroudja-Adamclisi (namely AdRom), and two from Greece (Elassona, Thessaly, namely Elassona 1 and Elassona 2) were processed at the University of Thessaly in Greece. DE samples were dried in an oven to 4.5% moisture content at 40 °C (Arnaud *et al.*, 2005). After drying, small pieces were grounded in a laboratory mill at full speed for 10 sec. Then, each sample powder was sieved through a standard sieve of 200 mesh (75 µm, Advantech Manufacturing Inc., New Berlin, WI, USA) and separately kept for 24 h in airproof jars. The commercially available DE formulations SilicoSec (Biofa GmbH, Münsingen, Germany) and PyriSec (Agrinova GmbH, Obrigheim/Mühleim, Germany) were also used. SilicoSec contains 92% SiO<sub>2</sub> while PyriSec contains 1.2% natural pyrethrum (25%), 3.1% piperonyl butoxide and 95.7% of SilicoSec (Vayias *et al.*, 2006).

In the 1<sup>st</sup> experiment untreated whole hard wheat (*Triticum durum* L., var. Miranda) and barley (*Hordeum vulgare* L., var. Laverda) was used in the tests. All DEs were applied in 1 kg of grain, at different amounts, creating lots with dose rates of 100, 300, 500 and 900 ppm (mg DE/kg of grain). The lots were placed in glass jars and shaken manually for 5 min to achieve equal distribution of the DE dust to the entire grain mass. For the 2<sup>nd</sup> experiment only one Greek (Elassona 1), one Romanian (PatRom) and one commercial DE (SilicoSec) were used. DEs were applied to whole hard wheat (var. Simeto), as previously, at doses of 300, 600 and 900 ppm. Additional 1 kg lots of wheat were sprayed with distilled water and used as control, for both experiments.

### Bioassays

In the 1<sup>st</sup> experiment, three samples of 50 g each, were taken from each lot, and placed in small cylindrical

glass vials (9 cm in diameter, 15 cm in height) with a different scoop that was inside each jar. The vials were covered with fine mesh for ventilation. Then, 30 adults were placed in each vial. The vials were then placed in controlled rooms set at 25 °C, 60% RH and continuous darkness. For the 2<sup>nd</sup> experiment, three samples of 20 g each, were taken from each lot, and placed in small plastic cylindrical vials (3 cm in diameter, 8 cm in height) with a different scoop that was inside each jar and 20 adults of each species were separately placed into each vial. The vials were covered with fine mesh as previously. The vials were then placed to incubators set at the combinations of three temperatures (20, 25 and 30 °C) and two RH levels (55 and 75%) in continuous darkness. Therefore, there were six temperature × RH combinations. In both experiments, the internal “necks” of the vials were covered by Fluon (Northern Products Inc., Woonsocket, USA), to prevent insects from escaping. Mortality of the exposed individuals in both experiments was assessed after 7, 14 and 21 days of exposure. All bioassays were carried out in three replicates, by preparing new lots.

### Data analysis


Adult control mortality was low (< 5%), therefore no correction was considered necessary (Abbott, 1925). Adult mortality was analyzed separately for each species and grain by using the Multivariate Analysis of Variance (MANOVA) Repeated Measures and Wilks' Lambda estimate, by using the JMP software (Sall *et al.*, 2001). In the 1<sup>st</sup> experiment, dose, DE and commodity were the main effects, while dose, DE, temperature and relative humidity were the main effects in the 2<sup>nd</sup> experiment. The associated interactions of the main effects were also included in the analysis. In both experiments, the repeated factor was exposure interval, while mortality was the response variable. Means were separated by the Tukey-Kramer honestly significant difference (HSD) test at 0.05 significance level (Sokal and Rohlf, 1995).

## Results

### 1<sup>st</sup> experiment

For *O. surinamensis* and *T. castaneum*, the main effects and the associate interactions between and within exposure intervals are presented in table 1. Mortality of *O. surinamensis* adults was 100% after 7 days of exposure at doses > 300 and 500 ppm of PyriSec treated wheat and barley respectively, while at the lowest dose mortality was 67% in treated wheat and 95.5% in treated barley (figure 1). For SilicoSec, after 21 days, mortality was 95 and 97% at the highest dose in barley and wheat respectively. In both commodities, the Greek (Elassona 1) and the Romanian (PatRom) DEs had the same performance with SilicoSec at 500 ppm. This was also evident at the lowest dose for all Greek and Romanian DEs but mortality did not exceed 42%. Generally, the tested DEs performed better in barley than in wheat.

In the case of *T. castaneum*, all adults were dead at 900 ppm in barley and wheat treated with PyriSec after 21 days of exposure. Generally, *T. castaneum* was less



**Figure 1.** Percent mortality (mean  $\pm$  SE) of *O. surinamensis* adults exposed to barley and wheat treated with six DEs (E1 = Ellassona 1, E2 = Ellassona 2, Pyri = PyriSec, R1 = PatRom, R2 = AdRom, Sil = SilicoSec) for 7, 14 and 21 days at four doses. Within each exposure, commodity and dose, means followed by the same letter are not significantly different. In all cases d.f. = 5, 53; Tukey-Kramer (HSD) test at  $p = 0.05$ . Where no letters exist, no significant differences were noted.

**Table 1.** Repeated-measures MANOVA parameters for main effects and associated interactions for mortality levels of *O. surinamensis* and *T. castaneum* adults between and within exposure intervals (for both species error d.f. = 384).

	d.f.	<i>O. surinamensis</i>		<i>T. castaneum</i>	
		F	p	F	p
<b>Between exposure intervals</b>					
Intercept	1	2595.9	< 0.01	981.7	< 0.01
DE	5	145.9	< 0.01	375.2	< 0.01
Commodity	1	22.7	< 0.01	44.0	< 0.01
Dose	3	66.3	< 0.01	117.4	< 0.01
DE × commodity	5	1.1	0.38	3.2	< 0.01
DE × dose	15	2.9	< 0.01	43.8	< 0.01
Commodity × dose	3	0.1	0.94	7.2	< 0.01
DE × commodity × dose	15	0.8	0.71	1.2	0.26
<b>Within exposure intervals</b>					
Exposure × DE	10	14.2	< 0.01	83.2	< 0.01
Exposure × commodity	2	6.9	< 0.01	21.1	< 0.01
Exposure × dose	6	9.3	< 0.01	28.9	< 0.01
Exposure × DE × commodity	10	2.2	< 0.01	5.0	< 0.01
Exposure × DE × dose	30	2.8	< 0.01	21.9	< 0.01
Exposure × commodity × dose	6	1.7	0.13	4.9	< 0.01
Exposure × DE × commodity × dose	30	1.2	0.19	2.9	< 0.01

**Table 2.** Repeated-measures MANOVA parameters for main effects and associated interactions for mortality levels of *S. oryzae* and *R. dominica* adults between and within exposure intervals (for both species error d.f. = 432).

	d.f.	<i>S. oryzae</i>		<i>R. dominica</i>	
		F	p	F	p
<b>Between exposure intervals</b>					
Intercept	1	11637.6	< 0.01	9222.2	< 0.01
DE	2	5.2	< 0.01	1.5	0.22
Dose	2	4216.5	< 0.01	4908.6	< 0.01
Temperature	2	539.3	< 0.01	510.0	< 0.01
RH	1	27.4	< 0.01	15.7	< 0.01
DE × dose	4	10.4	< 0.01	9.0	< 0.01
DE × temperature	4	1.5	0.21	1.7	0.14
DE × RH	2	0.2	0.83	0.1	0.94
Dose × temperature	4	211.7	< 0.01	281.6	< 0.01
Dose × RH	2	1.3	0.27	6.4	< 0.01
Temperature × RH	2	0.3	0.77	2.1	0.13
DE × dose × temperature	8	2.0	0.05	1.1	0.36
DE × dose × RH	4	0.1	0.99	0.1	0.97
DE × temperature × RH	4	0.1	0.99	0.1	0.98
Dose × temperature × RH	4	0.3	0.90	0.1	1.00
DE × dose × temperature × RH	8	0.1	1.00	0.3	0.97
<b>Within exposure intervals</b>					
Exposure × DE	4	8.2	< 0.01	6.5	< 0.01
Exposure × dose	4	120.6	< 0.01	202.0	< 0.01
Exposure × temperature	4	13.4	< 0.01	16.9	< 0.01
Exposure × RH	2	10.3	< 0.01	2.0	0.14
Exposure × DE × dose	8	2.1	0.04	1.4	0.18
Exposure × DE × temperature	8	3.5	< 0.01	3.2	< 0.01
Exposure × DE × RH	4	0.1	0.97	0.4	0.83
Exposure × dose × temperature	8	27.3	< 0.01	15.2	< 0.01
Exposure × dose × RH	4	0.3	0.89	0.4	0.78
Exposure × temperature × RH	4	0.2	0.95	0.3	0.89
Exposure × DE × dose × temperature	16	3.7	< 0.01	3.6	< 0.01
Exposure × DE × dose × RH	8	0.1	1.00	0.2	0.99
Exposure × DE × temperature × RH	8	0.1	1.00	0.2	0.99
Exposure × dose × temperature × RH	8	0.4	0.94	0.2	0.99
Exposure × DE × dose × temperature × RH	16	0.1	1.00	0.1	1.00

susceptible than *O. surinamensis* to the tested DEs. For SilicoSec adult mortality reached 74% only after 21 days at 900 ppm in wheat, while for the same exposure and dose in barley, mortality was 36%. Greek and Romanian DEs caused low mortality (< 25%) even after 21 days of exposure at the highest dose (figure 2). The local DEs and SilicoSec performed equally but for doses < 900 ppm. Higher mortality levels were recorded in wheat treated kernels compared with barley.

## 2<sup>nd</sup> experiment

For *S. oryzae* and *R. dominica*, the main effects and the associate interactions between and within exposure intervals are presented in table 2. Generally, *S. oryzae* adults were more susceptible than *R. dominica* adults to the tested DEs (figures 3, 4). Mortality levels for both species were high, but only at 900 ppm. After 21 days of exposure, mortality of *S. oryzae* and *R. dominica* adults reached 98% and 88% in the combination of 30 °C and 55% RH respectively. The increase of temperature, increased the efficacy of the tested DEs against adults of both species. This was more obvious at 900 ppm at both RH levels. For example, the Greek DE caused 58, 89 and 94% mortality to *S. oryzae* adults or 48, 79 and 85% to *R. dominica* adults at 20, 25 and 30 °C respectively at 55% RH after 21 days of exposure. The two RH levels influenced the insecticidal efficacy of DEs. Thus, higher mortality levels were recorded in the low RH for both species. Among the tested DEs there were no differences in their efficacy, with few exceptions. SilicoSec was more effective at 300 and 600 ppm at 20 °C compared to Greek and Romanian DEs, while at 900 ppm the Romanian DE was more effective compared with SilicoSec and Greek DE at 25 °C.

## Discussion


In the light of our findings, among DEs tested here, the most effective was PyriSec. The high mortality levels of *O. surinamensis* and *T. castaneum* adults recorded in the present study could be explained by the existence of natural pyrethrum and piperonyl butoxide in PyriSec through a possible synergism (Athanasassiou and Kavalieratos, 2005). Previous studies have also demonstrated the superiority of this DE upon other DEs. For example, Athanasassiou *et al.* (2004) reported that PyriSec caused significantly higher mortality of *S. oryzae* adults than the DEs Insecto and SilicoSec after 48 h of exposure in the treated substrate.

Of the insect species tested here, *O. surinamensis* was the most susceptible while *T. castaneum* the most tolerant to DEs, with the exception of PyriSec. *S. oryzae* and *R. dominica* were susceptible to the highest dose of all tested DEs but *R. dominica* was more tolerant than *S. oryzae*. Arthur (1991) reported that *O. surinamensis* adults were susceptible in wheat treated with the DE formulation Protect-It, given that both immediate and delay mortality were high. The susceptibility of this species to DEs could be attributed to the fact that it is mobile (Arthur, 1991) and thus, the probability of picking up DE particles is high. The mobility of an insect


species increases the susceptibility to DE (Rigaux *et al.*, 2001). Furthermore, *O. surinamensis* is a small insect, i.e., it can be dehydrated easily due to the DE activity (Korunic, 1998). Similarly, adults of *S. oryzae* are agile (Hill, 1990) while adults of *R. dominica* are moving slowly (Flinn and Hagstrum, 2011). *Tribolium* spp. are considered as the most tolerant stored-grain beetle species to DEs at the adult stage (Korunic, 1998; Fields and Korunic, 2000; Vayias and Athanasassiou, 2004; Athanasassiou *et al.*, 2011). This could be attributed to the fact that the DE particles are not easily attached on the body of *T. castaneum* (Fields and Korunic, 2000), given that this species is not hairy. Practically, the local DEs were ineffective against *T. castaneum* suggesting that longer exposure intervals and higher doses are required for the control of this species.

Temperature was associated positively with *R. dominica* and *S. oryzae* adult mortality and this was more obvious at the highest dose of all tested DEs. The increase of temperature increases insect mobility and the absorption of DE particles on the body surface. High temperatures also lead to increased respiration and water loss (Arthur, 2000b; Subramanyam and Roesli, 2000). However, the overall mortality of *R. dominica* adults did not exceed 49 or 44% at 20 °C and 55 or 75% RH respectively after 21 days of exposure. Similarly, the respective mortality values for *S. oryzae* were 59 and 54%. Thus, the tested DEs were not effective at 20 °C in both RH levels. This finding has been previously reported for a different DE (Athanasassiou *et al.*, 2014). The Romanian DE PatRom was highly efficacious mostly against *S. oryzae*, and in a lesser degree against *R. dominica*, at 900 ppm at 25 or 30 °C and 55% RH after 21 days of exposure. This finding is important because at 29.1 °C *S. oryzae* develops quickly (25 days) from egg to adult (Sharifi and Mills, 1971) and PatRom could offer a good level of protection to wheat against this pest. Furthermore, novel DEs with high insecticidal value in relatively low doses could be further improved with the addition of other substances, i.e., botanicals or insecticides which exhibit low mammalian toxicity, and consequently reduce the application dose (Athanasassiou and Korunic, 2007; Athanasassiou *et al.*, 2007; 2008; 2009; Vayias and Stephou, 2009). However, further processing and experimentation is needed to clarify this hypothesis.


Another important issue was the influence of RH to the Romanian DE PatRom. The high insecticidal efficacy of this DE against *S. oryzae* was slightly reduced with the increase of RH. When relative humidity is high, insects regulate water loss and thus their survival is increased in DE-treated substrate. Furthermore, at high air RH or grain moisture content levels, the DE particles become partially ineffective due the absorption of moisture (Subramanyam and Roesli, 2000; Mewis and Ulrichs, 2001). Newer studies have clearly demonstrated that even a small change of the level of relative humidity is able to influence the efficacy of DEs. For example, Athanasassiou *et al.* (2011) reported that mortalities of *R. dominica*, *S. oryzae* and *T. confusum* adults in wheat that had been previously treated with various DEs were lower at 65% than at 55%. Thus, the finding



**Figure 2.** Percent mortality (mean  $\pm$  SE) of *T. castaneum* adults exposed to barley and wheat treated with different DEs (E1 = Ellassona 1, E2 = Ellassona 2, Pyri = PyriSec, R1 = PatRom, R2 = AdRom, Sil = SilicoSec) for 7, 14 and 21 days at four doses. Within each exposure, commodity and dose, means followed by the same letter are not significantly different. In all cases d.f. = 5, 53; Tukey-Kramer (HSD) test at  $p = 0.05$ . Where no letters exist, no significant differences were noted.



**Figure 3.** Percent mortality (mean  $\pm$  SE) of *S. oryzae* adults exposed to wheat treated with three DEs (E1 = Ellassona 1, R1 = PatRom, Sil = SilicoSec) for 7, 14 and 21 days of under three temperatures and two RH levels. Within each RH, temperature and dose, means followed by the same letter are not significantly different. In all cases d.f. = 2, 26; Tukey-Kramer (HSD) test at  $p = 0.05$ . Where no letters exist, no significant differences were noted.



**Figure 4.** Percent mortality (mean  $\pm$  SE) of *R. dominica* adults exposed to wheat treated with three DEs (E1 = Elasona 1, R1 = PatRom, Sil = SilicoSec) for 7, 14 and 21 days of under three temperatures and two RH levels. Within each RH, temperature and dose, means followed by the same letter are not significantly different. In all cases d.f. = 2, 26; Tukey-Kramer (HSD) test at  $p = 0.05$ . Where no letters exist, no significant differences were noted.

of a DE which efficacy is stable to the increase of RH is particularly interesting, given that *S. oryzae* oviposition and survival of larvae are favored by elevated moisture content of the commodities (Singh *et al.*, 1974).

In the present study we found that the mortality levels of *O. surinamensis* and *T. castaneum* were different between barley and wheat. The fact that the type of grain determines the efficiency of DEs has been previously documented (Athanasioiu *et al.*, 2003; 2004, Kavallieratos *et al.*, 2005; 2010). Chiriloaie *et al.* (2014) also found that the efficacy of two Romanian and one Greek mined DEs was different among barley, maize, rice and wheat against *S. oryzae* adults. Our results are in agreement with these studies. Thus, for potential maximization of the efficiency of the management strategies, which are based on mined DEs, the type of commodity that is going to be treated and the target species are necessary to be known.

In summary, our results indicate that the Romanian DE PatRom tested here was found effective and compatible to the commercially available DEs, against *S. oryzae* adults mostly but also against *R. dominica* adults. However, further experimentation is needed on this DE against more stored-product insect species including studies on its influence against insects' progeny production in order to make a decision about the industrialization of this material.

## Acknowledgements

The study was supported by the project "Development of diatomaceous earth-based formulations from natural deposits from Greece and Romania, for stored-product protection against insects and mycotoxins" (General Secretary for Research and Technology, Greece, EPANII/ESPA, UEFISCDI Romania, Joint Research and Technology program between Greece and Romania).

## References

- ABBOTT W. S., 1925.- A method of computing the effectiveness of an insecticide.- *Journal of Economic Entomology*, 18: 265-267.
- ALDRYHIM Y. N., 1993.- Combination of classes of wheat and environmental factors affecting the efficacy of amorphous silica dust, dryacide, against *Rhizopertha dominica* (F.).- *Journal of Stored Products Research*, 29: 271-275.
- ANDRIĆ G. G., MARKOVIĆ M. M., ADAMOVIĆ M., ĐAKOVIĆ A., PRAZIĆ M. G., KLJAJIĆ P. J., 2012.- Insecticidal potential of natural zeolite and diatomaceous earth formulations against rice weevil (Coleoptera: Curculionidae) and red flour beetle (Coleoptera: Tenebrionidae).- *Journal of Economic Entomology*, 105: 670-678.
- ARNAUD L., HUONG T. T. L., BROSTAU X. Y., HAUBRUGE E., 2005.- Efficacy of diatomaceous earth formulations admixed with grain against populations of *Tribolium castaneum*.- *Journal of Stored Products Research*, 41: 121-130.
- ARTHUR F. H., 1991.- Immediate and delayed mortality of *Oryzaephilus surinamensis* (L.) exposed on wheat treated with diatomaceous earth: effects of temperature, relative humidity, and exposure interval.- *Journal of Stored Products Research*, 37: 13-21.
- ARTHUR F. H., 2000a.- Impact of food source on survival of red flour beetles and confused flour beetles (Coleoptera: Tenebrionidae) exposed to diatomaceous earth.- *Journal of Economic Entomology*, 93: 1347-1356.
- ARTHUR F. H., 2000b.- Toxicity of diatomaceous earth to red flour beetles and confused flour beetles (Coleoptera: Tenebrionidae): effects of temperature and relative humidity.- *Journal of Economic Entomology*, 93: 526-532.
- ARTHUR F. H., 2002.- Survival of *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) on wheat treated with diatomaceous earth: impact of biological and environmental parameters on product efficacy.- *Journal of Stored Products Research*, 38: 305-313.
- ARTHUR F. H., 2012.- Aerosols and contact insecticides as alternatives to methyl bromide in flour mills, processing plants, and food warehouses.- *Journal of Pest Science*, 85: 323-329.
- ARTHUR F. H., THRONE J. E., 2003.- Efficacy of diatomaceous earth to control internal infestations of rice weevil and maize weevil (Coleoptera: Curculionidae).- *Journal of Economic Entomology*, 96: 510-518.
- ATHANASSIOU C. G., KAVALLIERATOS N. G., 2005.- Insecticidal effect and adherence of PyriSec® in different grain commodities.- *Crop Protection*, 24: 703-710.
- ATHANASSIOU C. G., KORUNIC Z., 2007.- Evaluation of two new diatomaceous earth formulations, enhanced with abamectin and bitterbarkomycin, against four stored-grain beetle species.- *Journal of Stored Products Research*, 43: 468-473.
- ATHANASSIOU C. G., KAVALLIERATOS N. G., TSAGANOU F. C., VAYIAS B. J., DIMIZAS C. B., BUCHELOS C. T., 2003.- Effect of grain type on the insecticidal efficacy of SilicoSec against *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae).- *Crop Protection*, 22: 1141-1147.
- ATHANASSIOU C. G., KAVALLIERATOS N. G., ANDRIS N. S., 2004.- Insecticidal effect of three diatomaceous earth formulations against adults of *Sitophilus oryzae* (Coleoptera: Curculionidae) and *Tribolium confusum* (Coleoptera: Tenebrionidae) on oat, rye, and triticale.- *Journal of Economic Entomology*, 97: 2160-2167.
- ATHANASSIOU C. G., VAYIAS B. J., DIMIZAS C. B., KAVALLIERATOS N. G., PAPAGREGORIOU A. S., BUCHELOS C. T., 2005.- Insecticidal efficacy of diatomaceous earth against *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) and *Tribolium confusum* du Val (Coleoptera: Tenebrionidae) on stored wheat: influence of dose rate, temperature and exposure interval.- *Journal of Stored Products Research*, 41: 47-55.
- ATHANASSIOU C. G., KAVALLIERATOS N. G., PETEINATOS G. G., PETROU S. E., BOUKOUVALA M. C., TOMANOVIĆ Ž., 2007.- Influence of temperature and humidity on insecticidal effect of three diatomaceous earth formulations against larger grain borer (Coleoptera: Bostrychidae).- *Journal of Economic Entomology*, 100: 599-603.
- ATHANASSIOU C. G., KAVALLIERATOS N. G., VAYIAS B. J., STEPHOU V. K., 2008.- Evaluation of a new, enhanced diatomaceous earth formulation for use against the stored products pest, *Rhizopertha dominica* (Coleoptera: Bostrychidae).- *International Journal of Pest Management*, 54: 43-49.
- ATHANASSIOU C. G., KORUNIC Z., VAYIAS B. J., 2009.- Diatomaceous earths enhance the insecticidal effect of bitterbarkomycin against stored-grain insects.- *Crop Protection*, 28: 123-127.
- ATHANASSIOU C. G., KAVALLIERATOS N. G., VAYIAS B. J., TOMANOVIĆ Ž., PETROVIĆ A., ROZMAN V., ADLER C., KORUNIC Z., MILOVANOVIĆ D., 2011.- Laboratory evaluation of diatomaceous earth deposits mined from several locations in central and southeastern Europe as potential protectants against coleopteran grain pests.- *Crop Protection*, 30: 329-339.
- ATHANASSIOU C. G., KAVALLIERATOS N. G., LAZZARI F. A., 2014.- Insecticidal effect of Keepdry® for the control of *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) and *Rhizopertha dominica* (F.) (Coleoptera: Bostrychidae) on wheat under laboratory conditions.- *Journal of Stored Products Research*, 59: 133-139.

- BALDASSARI N., MARTINI A., 2014.- The efficacy of two diatomaceous earths on the mortality of *Rhyzopertha dominica* and *Sitophilus oryzae*.- *Bulletin of Insectology*, 67: 51-55.
- CHIRILOAIE A., CIOBANU M., DROSU S., 2013.- Preliminary studies regarding insecticidal effect of diatomaceous earth against the rice weevil, *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae), on stored wheat.- *Romanian Journal of Plant Protection*, 6: 27-30.
- CHIRILOAIE A., ATHANASSIOU C. G., VASSILAKOS T. N., FATU V., DROSU S., CIOBANU M., 2014.- Influence of grain type on the efficacy of some formulations of diatomaceous earth against the rice weevil (*Sitophilus oryzae* L.).- *Scientific Papers. Series A. Agronomy*, 57: 140-145.
- DESMARCHELIER J. M., DINES J. C., 1987.- Dryacide treatment of stored wheat: its efficacy against insects, and after processing.- *Australian Journal of Experimental Agriculture*, 27: 309-312.
- EBELING W., 1971.- Sorptive dusts for pest control.- *Annual Review of Entomology*, 16: 123-158.
- FIELDS P., KORUNIC Z., 2000.- The effect of grain moisture content and temperature on the efficacy of diatomaceous earths from different geographical locations against stored-product beetles.- *Journal of Stored Products Research*, 36: 1-13.
- FLINN P. W., HAGSTRUM D. W., 2011.- Movement of *Rhyzopertha dominica* in response to temperature gradients in stored wheat.- *Journal of Stored Products Research*, 47: 407-410.
- GOLOB P., 1997.- Current status and future perspectives for inert dusts for control of stored product insects.- *Journal of Stored Products Research*, 33: 69-79.
- HILL D. S., 1990.- *Pests of stored products and their control*.- Belhaven Press, London, UK.
- KAVALLIERATOS N. G., ATHANASSIOU C. G., VAYIAS B. J., BETSI P. C. C., 2010.- Insecticidal efficacy of fipronil against four stored-product insect pests: influence of commodity, dose, exposure interval, relative humidity and temperature.- *Pest Management Science*, 66: 640-649.
- KAVALLIERATOS N. G., ATHANASSIOU C. G., VAYIAS B. J., MAISTROU S. N., 2007.- Influence of temperature on susceptibility of *Tribolium confusum* (Coleoptera: Tenebrionidae) populations to three modified diatomaceous earth formulations.- *Florida Entomologist*, 90: 616-625.
- KAVALLIERATOS N. G., ATHANASSIOU C. G., PASHALIDOU F. G., ANDRIS N. S., TOMANOVIĆ Ž., 2005.- Influence of grain type on the insecticidal efficacy of two diatomaceous earth formulations against *Rhyzopertha dominica* (F.) (Coleoptera: Bostrychidae).- *Pest Management Science*, 61: 660-666.
- KAVALLIERATOS N. G., ATHANASSIOU C. G., VAYIAS B. J., KOTZAMANIDIS S., SYNODIS S. D., 2010.- Efficacy and adherence ratio of diatomaceous earth and spinosad in three wheat varieties against three stored product insect pests.- *Journal of Stored Products Research*, 46: 73-80.
- KORUNIC Z., 1998.- Diatomaceous earths, a group of natural insecticides.- *Journal of Stored Products Research*, 34: 87-97.
- LE PATOUREL G. N. J., 1986.- The effect of grain moisture content on the toxicity of a sorptive silica dust to four species of grain beetle.- *Journal of Stored Products Research*, 22: 63-69.
- MEWIS I., ULRICHS C., 2001.- Action of amorphous diatomaceous earth against different stages of the stored product pests *Tribolium confusum* (Coleoptera: Tenebrionidae), *Tenebrio molitor* (Coleoptera: Tenebrionidae), *Sitophilus granarius* (Coleoptera: Curculionidae) and *Plodia interpunctella* (Lepidoptera: Pyralidae).- *Journal of Stored Products Research*, 37: 153-164.
- NWAUBANI S. I., OPIT G. O., OTITODUM G. O., ADESIDA M. A., 2014.- Efficacy of two Nigeria-derived diatomaceous earths against *Sitophilus oryzae* (Coleoptera: Curculionidae) and *Rhyzopertha dominica* (Coleoptera: Bostrychidae) on wheat.- *Journal of Stored Products Research*, 59: 9-16.
- RIGAUX M., HAUBRUGE E., FIELDS P. G., 2001.- Mechanisms for tolerance to diatomaceous earth between strains of *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae).- *Entomologia Experimentalis et Applicata*, 101: 33-39.
- ROJHT H., ATHANASSIOU C. G., VAYIAS B. J., KAVALLIERATOS N. G., TOMANOVIĆ Ž., VIDRIH M., KOS K., TRDAN S., 2010.- The effect of diatomaceous earth of different origin, temperature and relative humidity against adults of rice weevil (*Sitophilus oryzae* [L.], Coleoptera: Curculionidae) in stored wheat.- *Acta Agriculturae Slovenica*, 95: 13-20.
- SALL J. E., LEHMAN A., CREIGHTON L., 2001.- *JMP Start Statistics. A guide to statistics and data analysis using JMP and JMP IN software*.- Duxbury Press, Belmont, CA, USA.
- SHARIFI S., MILLS R. B., 1971.- Developmental activities and behavior of rice weevil inside wheat kernels.- *Journal of Economic Entomology*, 64: 1114-1118.
- SINGH K., ARGAWAL N. S., GIRISH G. K., 1974.- The oviposition and development of *Sitophilus oryzae* (L.) in different high-yielding varieties of wheat.- *Journal of Stored Products Research*, 10: 105-111.
- SOKAL R. R., ROHLF F. J., 1995.- *Biometry. 3rd ed.*- Freeman, New York, USA.
- STATHERS T. E., DENNIF M., GOLOB P., 2004.- The efficacy and persistence of diatomaceous earth admixed with commodity against four tropical stored product beetle pests.- *Journal of Stored Products Research*, 40: 113-123.
- SUBRAMANYAM B., ROESLI R., 2000.- Inert dusts. pp. 321-380. In: *Alternatives to pesticides in stored-product IPM* (SUBRAMANYAM B., HAGSTRUM D. W., Eds).- Kluwer Academic Publishers, Dordrecht, The Netherlands.
- VAYIAS B. J., ATHANASSIOU C. G., 2004.- Factors affecting efficacy of the diatomaceous earth formulation SilicoSec against adults and larvae of the confused beetle *Tribolium confusum* du Val (Coleoptera: Tenebrionidae).- *Crop Protection*, 23: 565-573.
- VAYIAS B. J., STEPHOU V. K., 2009.- Factors affecting the insecticidal efficacy of an enhanced diatomaceous earth formulation against three stored-product insect species.- *Journal of Stored Products Research*, 45: 226-231.
- VAYIAS B. J., ATHANASSIOU C. G., KAVALLIERATOS N. G., TSESMELI C. D., BUCHELOS C. T., 2006.- Persistence and efficacy of two diatomaceous earth formulations and a mixture of diatomaceous earth with natural pyrethrum against *Tribolium confusum* Jacquelin du Val (Coleoptera: Tenebrionidae) on wheat and maize.- *Pest Management Science*, 62: 456-464.
- VAYIAS B. J., ATHANASSIOU C. G., KORUNIC Z., ROZMAN V., 2009.- Evaluation of natural diatomaceous earth deposits from south-eastern Europe for stored-grain protection: the effect of particle size.- *Pest Management Science*, 65:1118-1123.

**Authors' addresses:** Christos G. ATHANASSIOU, Thomas N. VASSILAKOS, Laboratory of Entomology and Agricultural Zoology, Department of Agriculture Crop Production and Rural Environment, University of Thessaly, Phytokou str., 38446 Nea Ionia, Magnissia, Greece; Nickolas G. KAVALLIERATOS (corresponding author: nick\_kaval@hotmail.com), Laboratory of Agricultural Zoology and Entomology, Department of Crop Science, Agricultural University of Athens, 75 Iera Odos str., 11855 Athens, Attica, Greece; Laboratory of Agricultural Entomology, Department of Entomology and Agricultural Zoology, Benaki Phytopathological Institute, 8 Stefanou Delta str., 14561 Kifissia, Attica, Greece; Andrei CHIRILOAIE, Viorel FĂTU, Sonica DROSU, Maria CIOBANU, Roxana DUDOIU, Research Development Institute for Plant Protection Bucharest, 8 Ion Ionescu de la Brad Blvd, District 1, 013813 Bucharest, Romania.

Received May 19, 2015. Accepted December 9, 2015.